



Case Report

Management of ranula with a combination of intra-lesional corticosteroid therapy and micro-marsupialization- A case report

Khushboo^{1,*}, Supreet Shirolkar¹, Monalisa Das¹, Subir Sarkar¹

¹Dept. of Pedodontic & Preventive Dentistry, Dr. R. Ahmed Dental College & Hospital, Kolkata, West Bengal, India



ARTICLE INFO

Article history:

Received 15-07-2020

Accepted 21-07-2020

Available online 24-10-2020

Keywords:

Ranula

Mucoceles

Intralesional corticosteroid

Micromarsupialization

ABSTRACT

Various forms of salivary gland swellings benign or malignant in nature are found in the oral cavity. An appropriate evaluation is needed for proper diagnosis and management of these lesions. Ranula, is one of the most common benign salivary gland lesions develops on the floor of the mouth. This paper presents a possible management of large sized ranula by a combination therapy of intra-lesional corticosteroid and Micro-marsupialization in 8 year old child according to the clinical condition.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

Mucous retention cysts or mucoceles are common oral benign lesions developed due to extravasation of the mucus from salivary glands where ranula is a comparatively rare form of mucocele found on floor of the mouth involving sublingual salivary gland. The name "ranula" is derived from the Latin word "rana" meaning "frog." The swelling of ranula resembles the transparent underbelly of a frog or air sacs. Normally the size of the lesion is greater than 2 cm and it appears as a painless, tense fluctuating dome-shaped vesicle, often with a blue hue.^{1,2} This happens unilaterally mostly lateral to oral cavity floor and is usually seen in young adults with similar male and female prevalence.³ Research shows the prevalence approximately 0.2 cases per 1000 people, and account for 6% of all oral sialo-cysts.⁴

In most cases the patho-physiology of ranula are not well understood and considered as unknown. The primary etiology for the development of ranula is partial obstruction of sublingual duct. This can contribute to the development of an epithelially-lined retention cyst, occurring in < 10% of all ranulas. The second most common factor is a

wound that causes significant damage to the duct or deeper regions of the sublingual gland's body, leading to mucus extravasations and a pseudocyst formation.⁵ The diagnosis, treatment planning and management of ranula is typically determined by a combination of history, clinical condition, histopathological appearance, and imaging studies. This paper presents management of large sized ranula in 8-year-old child with combination therapy of intra-lesional corticosteroid and micro-marsupialization.

2. Case Report

An 8 year old female patient reported to the department of Pedodontics and Preventive Dentistry with the chief complaint of painless swelling in the floor of the mouth with history of intermittent exaggeration and remission since past 4 months. General health of the patient was good with insignificant past dental, medical and family history. There was no history of trauma or injury related to oro-facial region. On intraoral examination, the lesion appeared as a translucent, dome-shaped bluish swelling in the right side of floor of mouth adjacent to the base of the tongue measuring approximately 1 cm in diameter. The lesion was smooth surfaced, fluctuating, soft in consistency and nontender

* Corresponding author.

E-mail address: khushboorana26@gmail.com (Khushboo).

on palpation. Reports of hematological and biochemical investigations were within normal limit. [Figure 1]

Based on history and clinical situation the patient was diagnosed with an oral ranula involving right sublingual gland. Evaluating the size of the lesion, age of the patient and patient's parents requirement a combination therapy of intralesional injection followed by micro-marsupialization (if required) was planned. On the first appointment, patient and parent consents were taken after thorough discussion of treatment plan along with related benefit and risk. In the next appointment, a topical anaesthetic was applied to cover the entire lesion for approximately 3 minutes and aspiration of mucus was done with 3 ml syringe. Then 1ml of betamethasone (Decdan, 4 mg/1 ml) was slowly injected into the base of the lesion followed by periphery with an insulin syringe (0.3×8 mm size, 31 gauge). The procedure was done gradually for being comfortable and painless. This procedure was repeated for 4 consecutive weeks (once in a week) until complete regression of the lesion. [Figure 2]

The patient was kept for 14 days observation and found recurrence of the lesion (5mm diameter in size). Next, the micro-marsupialization procedure was planned and discussed with the parent and appointment was fixed. Micro-marsupialization was performed on the next visit to drain the mucus and reduce the size of lesion. Under local anesthesia, a thick silk suture (3-0) was passed through the internal part of the lesion along its widest diameter followed by making a surgical knot and patient was called for follow up after 7 days. The 7th day healing was satisfactory without any post-surgical complications. Follow-up was done for 3 month, 6 months, patient was sound and showed no sign of recurrence. [Figures 3 and 4]

3. Discussion

Mucocele and ranulas located beneath the tongue are often confused with other entities. The broad differential diagnosis can include hemangioma, lymphangioma, dermoid cyst, benign or malignant salivary gland neoplasm, lipoma, abscess, venous lake, fibroma or benign mesenchymal neoplasm. The diagnosis is often depends on clinical appearance of the mass along with a typical history of healing and recurrence similar to the present case. Diagnosis of the ranula is often supported with investigations like high-resolution ultrasound or sonography which is able to assess the lesion up to 90% of benign versus malignant tumours.⁶ Ranula and mucocele can be demonstrated more clearly through the use of colour Doppler imaging. Computed tomography or magnetic resonance imaging are seldom necessary except in the case of a large or 'plunging' ranula, which may breach the mylohyoid muscle and present in the lateral aspect of the neck.⁷ Detailed imaging can define the extent of the lesion and may be critical for the planning of surgery. Computed tomography and magnetic resonance imaging do




Fig. 1: Pre-operative photograph-Intraoral clinical appearance of lesion.




Fig. 2: A) Mucous aspiration by 18-gauge needle and syringe B) 1ml of Betamethasone injection and 31 gauge insulin syringe.

not differentiate between benign and malignant processes. Thus, a biopsy may be required for this purpose. The histological difference between ranula and mucocele is, the ranula is lined by granulation tissue, consisting of swelling and congested vasculature whereas, mucocele consists of ductal epithelium.⁸

Managing a ranula depends on a variety of variables but primarily on its size and location. In the process of removing stones from the hilum of the submandibular gland, a technique has been developed which requires that the sublingual gland be mobilized and rotated laterally on the




Fig. 3: A) recurrent lesion after intralesional corticosteroid therapy. B) post surgical photograph of micro-marsupialization of lesion.



Fig. 4: Seven days post operative photograph.

floor of the mouth.⁹ Small, well-located ranulas respond to local excision much the same as mucoceles on the lower lip. The surgical removal of ranula along with the associated sublingual gland is the most predictable method of getting rid of the lesion. Although, it is not necessarily an easy operation, but can be considered more successful if carried out in a clean, bloodless environment. Unfortunately, the morbidity of the removal of sublingual gland is relatively high.

Also, the chance of injury to the Wharton duct (2%), bleeding (1% to 2%), infection (1% to 2%) or lingual nerve paraesthesia (2% to 12%) may increase. A number of more progressive approaches that are consistent with office practices have been suggested, including laser de-roofing,¹⁰ cryotherapy, suture ligation, and marsupialization variation.¹¹

Micromarsupialization has been shown to be a simple, fairly non-invasive, painless, efficient and low recurrence technique used by Amaral et al.(2012) and Sagari et al(2012) to treat oral ranulas and selected mucoceles, in which all cases showed complete healing within 30 days of the procedure. Nonetheless, these surgical procedures have many drawbacks, such as trauma, discomfort, lip disfigurement, damage to adjacent vital structures, and ducts

that contribute to the creation of satellite lesions, and can also be costly for the patient.¹² Thus, a non-surgical treatment regimen with extremely active corticosteroids (betamethasone) has been introduced. Intracystic injection therapy with OK-432 (a lyophilized mixture of low virulence strain of streptococcus pyogenes incubated with benzyl penicillin) has been reported to be relatively safe and can be used as a surgical substitute for ranula therapy.¹³ Ranula can be treated with a silk suture placed in the cyst dome as a process of micro-marsupialization. Pandit and Park(2002) suggested that successful pediatric oral cavity management may involve observation of spontaneous resolution for 5 months. Unless the lesion doesn't regularly heal or recur, surgical therapy is prescribed. Hence the decision to perform combination therapy of intralesional corticosteroid and micro-marsupialization was taken as per the clinical situation.^{14,15}

4. Conclusion

Surgical management of oral lesion is often difficult in pediatric patients under local anaesthesia. Therefore, the combination therapy of intralesional corticosteroid and micro-marsupialization can be a treatment option instead of direct surgical excision in children. More cases with long-term observations are needed to assess the success of this therapy in children.

5. Source of Funding

None.

6. Conflict of Interest

None.

References

1. Fagan J. Ranula and sublingual salivary gland excision. *Open Access Atlas Otolaryngol.* 2012;.
2. Axell T. A prevalence study of oral mucosal lesions in an adult Swedish population". *Odontologisk Revy.* 1976;36:1–103.
3. Kokong D, Iduh A, Chukwu I, Mugu J, Nuhu S, Augustine S, et al. Ranula: Current Concept of Pathophysiologic Basis and Surgical Management Options. *World J Surg.* 2017;41(6):1476–81.
4. Olojede ACO, Ogundana OM, Emeka CI, Adewole RA, Emmanuel MM, Gbotolorun OM, et al. Plunging ranula: surgical management of case series and the literature review. *Clin Case Rep.* 2018;6(1):109–14.
5. Chauhan R, Chauhan V, Lele G, Shirol D. Ranula in an adolescent patient. *J Dent Allied Sci.* 2014;3(2):105.
6. Kessler AT, Bhatt AA. Review of the Major and Minor Salivary Glands, Part 2: Neoplasms and Tumor-like Lesions. *J Clin Imaging Sci.* 2018;8:48.
7. Sheikh M. Plunging ranula of the submandibular area. *Dent Res J.* 2011;8(1):114–8.
8. Boulos M, Cheng A. Case 1: What is that in your mouth?. *Paediatr Child Health.* 2006;11:107–8.
9. Leppi TJ. Gross Anatomical Relationships Between Primate Submandibular and Sublingual Salivary Glands. *J Dent Res.* 1967;46(2):359–65.

10. Barak S, Horowitz I, Katz J, Kaplan I. Experiences with the CO₂ Laser in the Surgical Treatment of Intraoral Salivary Gland Pathology. *J Clin Laser Med Surg*. 1991;9(4):295–9.
11. Fukase S, Inamura K, Ohta N, Aoyagi M. Treatment of Ranula with Intracystic Injection of the Streptococcal Preparation Ok-432. *Ann Otol, Rhinol Laryngol*. 2003;112(3):214–20.
12. Piazzetta CM, Torres-Pereira C, Amenábar JM. Micro-marsupialization as an alternative treatment for mucocele in pediatric dentistry. *Int J Paediatr Dent*. 2012;22(5):318–23.
13. Yoshizawa K, Moroi A, Kawashiri S, Ueki K. A Case of Sublingual Ranula That Responded Successfully to Localized Injection Treatment with OK-432 after Healing from Drug Induced Hypersensitivity Syndrome. *Case Rep Dent*. 2016;2016:1–5.
14. Pandit RT, Park AH. Management of Pediatric Ranula. *Otolaryngology–Head Neck Surg*. 2002;127(1):115–8.
15. Harrison JD. Modern management and patho physiology of ranula: Literature review. *Head Neck*. 2010;32:1310–20.

Author biography

Khushboo 2nd Year Post Graduate

Supreet Shirolkar 1st Year Post Graduate

Monalisa Das Assistant Professor

Subir Sarkar Professor and HOD

Cite this article: Khushboo, Shirolkar S, Das M, Sarkar S. **Management of ranula with a combination of intra-lesional corticosteroid therapy and micro-marsupialization- A case report.** *Int Dent J Students Res* 2020;8(3):114-117.